




2015 내시경 퀴즈 (#1-#5)

성균관대학교 의과대학 내과 이준행

Q1. 소견과 진단은?




A1. BGU active, Hp (+)




After PPI and
H. pylori eradication

Q2. 소견과 진단은?




A2. BGU active 2, Hp (+)



After PPI and
H. pylori eradication

Q3. 소견과 진단은?




A3. EGC

Early gastric carcinoma

1. Location : lower third, Center at antrum and greater curvature
2. Gross type : EGC type IIc
3. Histologic type : **tubular adenocarcinoma, poorly differentiated**
4. Histologic type by Lauren : diffuse
5. Size : 3.2x1.6 cm
6. Depth of invasion : **invades submucosa (sm3)** (pT1b)
7. Resection margin: free from carcinoma, safety margin: proximal 7.3 cm, distal 5.3 cm
8. Lymph node metastasis : **metastasis to 6 out of 47 regional lymph nodes (pN2)**
(perinodal extension: absent)
9. Lymphatic invasion : present
10. Venous invasion : present(intramural)
11. Perineural invasion : not identified
12. Peritoneal cytology : negative
13. AJCC stage by 7th edition: T1b N2

Q4. 소견과 진단은?




A4. AGC



Advanced gastric carcinoma

1. Location : upper third,
2. Gross type : Borrmann type 3
3. Histologic type : **tubular adenocarcinoma, poorly differentiated**
4. Histologic type by Lauren : diffuse
5. Size : 3.3x2 cm
6. Depth of invasion : **penetrates subserosal connective tissue (pT3)**
7. Resection margin: free from carcinoma
8. Lymph node metastasis : **metastasis to 13 out of 44 regional lymph nodes (pN3a)**
9. Lymphatic invasion : present
10. Venous invasion : not identified
11. Perineural invasion : present
15. Peritoneal cytology : negative

Q5. 소견과 진단은?



A5. BGU with *Candida*



After PPI (4 weeks)
plus fluconazole (1 week)



Acting 선생님은 PDF로 되어있는 문제를 모두에게 전달해 주세요.
각자 소견과 진단(first impression 하나만)을 A4 용지에 작성하여
다음 주 월요일 오후까지 acting 선생님께 전달해 주세요.
Acting 선생님은 목요일 집담회때 발표해 주세요.